

NEW LOWER BOUNDS FOR THE SIZE OF A NON-TRIVIAL LOOP IN THE COLLATZ $3x+1$ AND GENERALIZED $Px+Q$ PROBLEM

ROUPAM GHOSH

Abstract

In the Collatz $3x+1$ problem, there are 3 possibilities: Starting from any positive number, we either reach the trivial loop $(1,4,2)$, end up in a non-trivial loop, or go until infinity. In this paper, we shall show that if a non-trivial loop with m odd numbers exists, then its minimum odd number is bounded above by a function of m . We shall also use that bound to calculate the least number of odd elements required for a non-trivial loop to exist. Also, the generalized bounds for the $px+q$ problem are given.

Introduction to the Collatz problem

Consider the function,

$$f(x) = \begin{cases} \frac{x}{2} & \text{if } x \text{ is even} \\ 3x + 1 & \text{if } x \text{ is odd} \end{cases}$$

The Collatz $3x+1$ conjecture states that for every positive integer x , there exists an integer $d(\geq 0)$ corresponding to x such that $f^{(d)}(x) = 1$, where $f^{(d)}(x) = f(f(f(\dots d \text{ times} \dots (f(f(f(x))\dots)))$). For example, the iterations of f on a few numbers are given below:

$$(1) \quad \begin{aligned} &1 \rightarrow 1 \\ &2 \rightarrow 1 \\ &3 \rightarrow 10 \rightarrow 5 \rightarrow 16 \rightarrow 8 \rightarrow 4 \rightarrow 2 \rightarrow 1 \\ &4 \rightarrow 2 \rightarrow 1 \\ &5 \rightarrow 16 \rightarrow 8 \rightarrow 4 \rightarrow 2 \rightarrow 1 \\ &6 \rightarrow 3 \rightarrow 10 \rightarrow 5 \rightarrow 16 \rightarrow 8 \rightarrow 4 \rightarrow 2 \rightarrow 1 \\ &\text{and so on...} \end{aligned}$$

Considering only odd numbers

Now, let us consider only odd integers of the Collatz sequence and modify the function f to a function T defined on odd integers such that $T(x) = \frac{(3x+1)}{2^k}$, where k is the highest power of two in which 2^k divides $3x + 1$.

For example, we have for $x = 7$:

$$(2) \quad T(7) = 11, T^{(2)}(7) = 17, T^{(3)}(7) = 13, T^{(4)}(7) = 5, T^{(5)}(7) = 1, T^{(6)}(7) = 1, T^{(7)}(7) = 1, \dots$$

$$k_1 = 1, k_2 = 1, k_3 = 2, k_4 = 3, k_5 = 4, k_6 = 2, k_7 = 2, \dots$$

and so on, where k_i is the highest power of two which divides $3T^{(i-1)}(x) + 1$. The iterations of T on a few odd numbers are given below:

$$(3) \quad \begin{aligned} &1 \rightarrow 1 \\ &3 \rightarrow 5 \rightarrow 1 \\ &5 \rightarrow 1 \\ &7 \rightarrow 11 \rightarrow 17 \rightarrow 13 \rightarrow 5 \rightarrow 1 \\ &9 \rightarrow 7 \rightarrow 11 \rightarrow 17 \rightarrow 13 \rightarrow 5 \rightarrow 1 \\ &11 \rightarrow 17 \rightarrow 13 \rightarrow 5 \rightarrow 1 \\ &13 \rightarrow 5 \rightarrow 1 \\ &15 \rightarrow 23 \rightarrow 35 \rightarrow 53 \rightarrow 5 \rightarrow 1 \\ &\text{and so on...} \end{aligned}$$

Some inequalities

Starting with any arbitrary odd integer a_1 , let $a_r = T^{(r)}(a_1)$. Let us suppose a non-trivial loop exists and consists of m odd numbers. Let us consider $2_1^k, 2_2^k, \dots, 2_m^k$ to be the highest powers of two which divide $3a_1 + 1, 3a_2 + 1, \dots, 3a_m + 1$, respectively.

And let us denote $S_r = k_1 + k_2 + \dots + k_r$. Then we shall have

$$(4) \quad a_{m+1} = \frac{3^m}{2^{S_m}} a_1 + \frac{c_m}{2^{S_m}}$$

where $c_m = 3^{m-1} + 3^{m-2}2^{S_1} + \dots + 3^2 2^{S_{m-3}} + 3^1 2^{S_{m-2}} + 2^{S_{m-1}}$.

Therefore, by definition,

$$(5) \quad \begin{aligned} 2^{S_r} &= 2^{k_1+k_2+\dots+k_r} \\ &= \frac{(3a_1+1)}{a_2} \frac{(3a_2+1)}{a_3} \dots \frac{(3a_r+1)}{a_{r+1}} \\ &= \left(3 + \frac{1}{a_1}\right) \left(3 + \frac{1}{a_2}\right) \dots \left(3 + \frac{1}{a_r}\right) \frac{a_1}{a_{r+1}} \end{aligned}$$

Then for $a_1 = a_{m+1}$, i.e., for a loop containing m odd numbers, we shall have

$$(6) \quad 2^{S_m} = \left(3 + \frac{1}{a_1}\right) \left(3 + \frac{1}{a_2}\right) \dots \left(3 + \frac{1}{a_m}\right)$$

Let us consider a_{min} to be the minimum among $a_1, a_2, \dots, a_m$. Then we have

$$(7) \quad \begin{aligned} 3^m &< 2^{S_m} < \left(3 + \frac{1}{a_{min}}\right)^m \\ m(\log_2 3) &< S_m < m(\log_2 3) + m \log_2 \left(1 + \frac{1}{3a_{min}}\right) \end{aligned}$$

Deriving the bound

Now, since S_m is a positive integer, if

$$[m(\log_2 3) + m \log_2(1 + \frac{1}{3a_{min}})] - [m(\log_2 3)] = 0$$

then no integer solution S_m exists (where $[x]$ is the floor function), and this will imply that no loop exists with m-odd numbers. Here, $\{x\}$ denotes $x - [x]$, the fractional part of x . Now, if the above condition is true then we have

$$m(\log_2 3) + m \log_2(1 + \frac{1}{3a_{min}}) - \{m(\log_2 3) + m \log_2(1 + \frac{1}{3a_{min}})\} = m(\log_2 3) - \{m(\log_2 3)\}$$

$$\text{or } m \log_2(1 + \frac{1}{3a_{min}}) = \{m(\log_2 3) + m \log_2(1 + \frac{1}{3a_{min}})\} - \{m(\log_2 3)\}$$

So if there is an integer solution for S_m , we must have

$$m \log_2(1 + \frac{1}{3a_{min}}) > \{m(\log_2 3) + m \log_2(1 + \frac{1}{3a_{min}})\} - \{m(\log_2 3)\}.$$

This leaves us with only two possibilities, $m \log_2(1 + \frac{1}{3a_{min}}) > 1$ or $m \log_2(1 + \frac{1}{3a_{min}}) < 1$.

Bound for the case $m \log_2(1 + \frac{1}{3a_{min}}) > 1$:

$$m \log_2(1 + \frac{1}{3a_{min}}) > 1$$

Rearranging the terms of $m \log_2(1 + \frac{1}{3a_{min}}) > 1$, we get

$$a_{min} < \frac{1}{3(2^{\frac{1}{m}} - 1)}$$

We shall define $\alpha(m)$ to be $\frac{1}{3(2^{\frac{1}{m}} - 1)}$.

Bound for the case $m \log_2(1 + \frac{1}{3a_{min}}) < 1$:

From above, if a solution for S_m exists then we have

$$m \log_2(1 + \frac{1}{3a_{min}}) > \{m(\log_2 3) + m \log_2(1 + \frac{1}{3a_{min}})\} - \{m(\log_2 3)\}$$

But, since $m \log_2(1 + \frac{1}{3a_{min}}) < 1$, $\{m \log_2(1 + \frac{1}{3a_{min}})\} = m \log_2(1 + \frac{1}{3a_{min}})$.

Hence,

$$\{m \log_2(1 + \frac{1}{3a_{min}})\} > \{m(\log_2 3) + m \log_2(1 + \frac{1}{3a_{min}})\} - \{m(\log_2 3)\}$$

or

$$\{m(\log_2 3)\} + \{m \log_2(1 + \frac{1}{3a_{min}})\} > \{m(\log_2 3) + m \log_2(1 + \frac{1}{3a_{min}})\}$$

This is only possible if

$$\{m(\log_2 3)\} + \{m \log_2(1 + \frac{1}{3a_{min}})\} > 1.$$

Therefore, we have

$$m \log_2(1 + \frac{1}{3a_{min}}) > 1 - \{m(\log_2 3)\}.$$

Rearranging the terms, we get

$$a_{min} < \frac{1}{3(2^{\frac{1-\{m \log_2 3\}}{m}} - 1)}.$$

We shall define $\beta(m)$ to be $\frac{1}{3(2^{\frac{1-\{m \log_2 3\}}{m}} - 1)}$.

Some new definitions:

We shall divide the possible non-trivial loops of a Collatz sequence into two classes: We call loops satisfying the equation

$$m \log_2(1 + \frac{1}{3a_{min}}) > 1$$

α -loops and loops satisfying the equation

$$m \log_2(1 + \frac{1}{3a_{min}}) < 1$$

β -loops.

Note that for α -loops we have $a_{min} < \alpha(m)$ and for β -loops we have $a_{min} < \beta(m)$.

Computing the least number of odd numbers a loop must have

The Collatz algorithm has been tested and found to always reach 1 for all numbers $\leq 19 \times 2^{58}$ (Oliveira e Silva 2008).

For α -loops, we have

$$a_{min} < \frac{1}{3(2^{\frac{1}{m}} - 1)}$$

i.e.,

$$m > \frac{\log(2)}{\log(1 + \frac{1}{3a_{min}})}$$

Putting $a_{min} = 19 \times 2^{58} - 1$, we get $m > 11387806137299329586$.

Hence, if an α -loop exists we must have at least 11, 387, 806, 137, 299, 329, 586 odd numbers in that loop.

For β -loops we have

$$\frac{m}{1 - \{m \log_2 3\}} > \frac{\log(2)}{\log(1 + \frac{1}{3a_{min}})}$$

Given below is PARI/GP code I used for calculating the value of m for β -loops.

```
lhs(x) = ( x / ( 1 - frac( x*log(3.0)/log(2.0) ) ) );
rhs(x) = ( log(2)/log(1 + 1/(3*x) ) );
{
    a = rhs( 19 * ( 2^58) - 1 );
    m = 2;
    while(1,
        if( lhs(m) > a,
            print(m," is our desired result ");
            quit;
        ,
            print(m, " has been checked...");
        );
    m++;
};
}
```

The value for β -loop is 6,586,818,670.

That is, at least 6,586,818,670 odd numbers are required to form a β -loop, which is much larger than the current largest known lower bound for the length of a non-trivial cycle (Sinisalo, 2003).

Bounds for the generalized $px+q$ problem

Consider the function,

$$f(x) = \begin{cases} \frac{x}{2} & \text{if } x \text{ is even} \\ px + q & \text{if } x \text{ is odd. Here } p \text{ and } q \text{ are both positive odd numbers} \end{cases}$$

Then we shall have

$$\alpha(m) = \frac{q}{p(2^{\frac{1}{m}} - 1)}$$

$$\beta(m) = \frac{q}{p(2^{\frac{1 - \{m \log_2 p\}}{m}} - 1)}$$

We denote this fact by introducing this new notation,

$$\alpha_m(p, q) = \frac{q}{p(2^{\frac{1}{m}} - 1)}$$

$$\beta_m(p, q) = \frac{q}{p(2^{\frac{1 - \{m \log_2 p\}}{m}} - 1)}$$

For example, in our case of $3x + 1$, we have

$$\alpha_m(3, 1) = \frac{1}{3(2^{\frac{1}{m}} - 1)}$$

Acknowledgements

I thank Craig Alan Feinstein for giving me feedback on various important steps, providing me useful insights, and also amongst other things, for being a good friend. Also, my thanks goes to Eric Farin for helping me by running the program on his computer. I thank my mom, sister, and dad for making my life wonderful.

References

- (1) Weisstein, Eric W. "Collatz Problem." From MathWorld—A Wolfram Web Resource. <http://mathworld.wolfram.com/CollatzProblem.html>
- (2) Toms Oliveira e Silva, <http://www.ieeta.pt/~tos/3x+1.html>
- (3) Shalom Eliahou (1993), "The $3x+1$ Problem, New Lower Bounds on a Nontrivial Cycle Lengths", Discrete Math 118
- (4) Busido Chisale (1994), "Cycles in Collatz Sequences" Publ. Math. Debrecen 45
- (5) Lorentz Halbeisen and Norbert Hungerbühler (1997), "Optimal Bounds for the Length of Rationnal Collatz Cycles", Acta Arithmetica 78
- (6) Matti Sinisalo (2003), "On the Minimal Cycle Length of the Collatz Sequences", Univ of Oulu, Finland
- (7) The PARI Group - PARI/GP version 2.3.4, <http://pari.math.u-bordeaux.fr/>